

(REVIEW ARTICLE)



Advances in understanding the cellular composition and molecular signatures of the adult *Drosophila* eye through single-cell RNA sequencing

Blessie Pradeeka Gudelly¹ and Komal K. B. Raja^{2,*}

¹ Osmania Medical College, Department of Medicine, Hyderabad, Telangana, 500095, India.

² Department of Biotechnology, Malla Reddy Engineering College, Medchal, Telangana, 500100, India.

Magna Scientia Advanced Research and Reviews, 2024, 11(02), 365–369

Publication history: Received on 23 June 2024; revised on 02 August 2024; accepted on 05 August 2024

Article DOI: <https://doi.org/10.30574/msarr.2024.11.2.0130>

Abstract

The fruit fly *Drosophila melanogaster* has long served as a model organism for studying genetic and developmental processes. The *Drosophila* eye, with its well-characterized structure and genetic tractability, is a prime system for exploring the principles of differentiation, cell type specification, and neurogenesis. Recent advances in single-cell RNA sequencing (scRNA-seq) technology have revolutionized our ability to study gene expression at the individual cell level, offering unprecedented insights into cellular heterogeneity within complex tissues. This review highlights the application of scRNA-seq to the adult *Drosophila* eye, emphasizing its role in delineating the transcriptomic landscape of its diverse cell types. By profiling individual cells, researchers have identified novel marker genes for all major cell types in the eye, providing a detailed cellular atlas and uncovering previously unknown cellular subtypes. The review also discusses the broader implications of these methodologies for developmental biology, the importance of functional validation of identified markers, and the potential of integrating scRNA-seq data with other omics approaches. Overall, single-cell genomics of the *Drosophila* eye sets a new standard for cellular resolution, offering valuable insights into the complexities of gene regulation and cellular diversity, with significant implications for understanding development and disease.

Keywords: Single cell RNA-seq; Bioinformatics; *Drosophila* eye; Single cell omics

1. Introduction

Understanding the principles underlying development, differentiation, and pattern formation is central to developmental biology. For over a century, the fruit fly *Drosophila melanogaster* has been an exemplary model for uncovering the genetic basis of development and disease [1-5]. *Drosophila* organs and tissues are extensively utilized to investigate the genes, pathways, and signaling mechanisms involved in development and disease. The *Drosophila* eye, in particular, has served as a robust system for exploring differentiation, planar cell polarity, cell type specification, cell death, survival, and neurogenesis[1].

The adult *Drosophila* eye comprises approximately 800 light-sensing units called ommatidia, each containing eight photoreceptor neurons and several accessory cells. The ease of genetic manipulation in the laboratory, availability of numerous genetic tools, and high conservation with human genes make the *Drosophila* eye an excellent model for studying gene function and genes involved in neurodegenerative diseases. The *Drosophila* eye has been instrumental in screening genes within unidentified disease networks.

Previous studies have extensively characterized the developmental processes and genetic pathways involved in eye formation[6]. However, the precise molecular identity and functional heterogeneity of the various cell types within the

* Corresponding author: Komal K. B. Raja

mature adult eye have remained less well-defined. Data generated from pooled cell populations, such as bulk RNA-seq technologies, lose cell type information and provide an average gene expression measurement across all cells in the sample. This masks differences between individual cells in a given population and often overlooks rare subpopulations or unknown cell types that could be drivers of disease or development. Thus, single-cell approaches are necessary for studying tissues, particularly important model systems like *Drosophila* organs and tissues.

1.1. The Advent of Single Cell RNA Sequencing

The advent of single-cell RNA sequencing (scRNA-seq) technology has revolutionized our ability to delineate the transcriptomic landscape of individual cells within any tissue. This technology offers a detailed understanding of cell type-specific gene expression patterns and uncovers previously unknown cellular subtypes. scRNA-seq is a powerful tool for dissecting cellular heterogeneity in complex tissues [7, 8]. The adult *Drosophila* eye, a well-established model for studying development and neurogenetics, presents a unique opportunity to apply scRNA-seq to uncover the cellular diversity and molecular signatures of its constituent cells [6, 9].

1.2. Advances in Dissociation Methods and High-Resolution Data Generation

Recent single-cell studies on the adult *Drosophila* eye have utilized the 10X Genomics platform. For instance, one study generated a single-cell nuclei sequence resource for the entire adult *Drosophila* fly, including the eye [10]. However, the eye data showed that the transcriptomes of neuronal and non-neuronal cells were mixed, confounding the results. Moreover, this data was based on single nuclei, potentially missing transcripts present in whole cells. Therefore, the resolution of data from adult nuclei was insufficient to provide a detailed cellular landscape of the adult eye. This limitation was expected, given the challenge of generating live single cells from the adult *Drosophila* eye, where cells are tightly held together in a crystalline lattice.

In contrast, the study by Yeung et al. (2022) [11], titled "Single cell RNA sequencing of the adult *Drosophila* eye reveals distinct clusters and novel marker genes for all major cell types," significantly advanced dissociation methods and generated data from whole cells from the adult *Drosophila* eye. This study profiled cells from all cell types in the adult eye, representing a significant advancement in understanding the cellular composition and molecular signatures of the adult *Drosophila* eye. The research leveraged the power of scRNA-seq to provide a comprehensive and high-resolution map of gene expression profiles of individual cells in the adult eye, offering new insights into the complexity of this essential organ.

2. Innovative Use of scRNA-seq in Detailing Cellular Landscape

The application of scRNA-seq to the adult *Drosophila* eye is both innovative and impactful. This technique allows for the dissection of gene expression at an unprecedented resolution, revealing distinct cellular clusters that traditional bulk RNA sequencing methods might overlook [12]. Previous studies on *Drosophila* pigmentation patterns have highlighted the importance of understanding gene regulation at a detailed and single-cell level, which this research achieves in the eye through scRNA-seq [13, 14].

2.1. Identification of Novel Marker Genes

One of the standout achievements in recent research is the identification of novel marker genes for all major cell types in the *Drosophila* eye, which was not possible in previous studies due to lower resolution in profiling single cells. These novel markers are invaluable for distinguishing between different cell types and understanding their unique functions and developmental pathways [15]. For example, the gene *CG2082* is specifically expressed in R8 neurons. While little is known about the function of this gene in R8 neurons, its discovery opens new avenues for neuronal research.

2.2. Comprehensive Cellular Atlas

This research also provides a detailed cellular atlas of the *Drosophila* eye, highlighting the diversity and complexity of cell types present. This comprehensive map is a crucial resource for researchers studying eye development, function, and disease. Similar comprehensive approaches have been applied to other *Drosophila* tissues, such as the larval eye, providing insights into photoreceptor development [16, 17]. Integrating the adult eye data with larval eye data will provide insights into differentiation and pattern formation during development.

2.3. Broad Implications for Developmental Biology

Beyond the *Drosophila* eye, the methodologies and findings from single-cell RNA sequencing studies have broad implications for developmental biology. The approach can be adapted to other tissues and model organisms, facilitating

a deeper understanding of cellular heterogeneity and gene regulation across different biological systems[18]. For instance, since the *Drosophila* eye has a cuticle, the single-cell dissociation technique can be utilized to isolate single cells from other tissues with a hardened cuticle. This adaptability is evident in the use of scRNA-seq to study developmental timelines and novel markers in the *Drosophila* larval eye [16, 17, 19]

3. Limitations and Future Directions

3.1. Functional Validation of Marker Genes and Comparative Analyses

While the identification of novel marker genes is a significant contribution, further functional validation of these markers would strengthen the findings. Sometimes a gene may be expressed but may not have any function. Understanding the roles these genes play in cell function and development would provide deeper insights into their biological significance. Functional studies could draw on techniques such as RNA in situ hybridization and imaging techniques to validate gene expression patterns, as demonstrated in previous work on *Drosophila* species [20-23]. Additionally, well-established genomic tools in *Drosophila* such as CRISPR, the FRT-FLP system, and the UAS-GAL4 system can be used to mutate or misexpress genes in any tissue and at any developmental stage. Using these methods, the function of these genes can be elucidated.

Further, a comparative analysis with scRNA-seq data from other model organisms or different developmental stages of the *Drosophila* eye could offer additional context and enhance the study's impact. Such comparisons could reveal conserved and divergent gene expression patterns, contributing to a broader understanding of eye development and evolution [24]. Comparative studies have already proven valuable in understanding gene regulation in the context of phylogeny and systematics [25].

3.2. Integration with Other Omics Data

Integrating the scRNA-seq data with other omics approaches, such as proteomics or epigenomics, could provide a more holistic view of the molecular mechanisms governing cell identity and function in the *Drosophila* eye. This multi-omics approach would enrich the findings and potentially uncover new regulatory networks [26]. For instance, integrative analyses have identified cell type-specific targets of the transcription factor Pointed in the *Drosophila* eye, highlighting the utility of such approaches [19].

4. Conclusion

In summary, scRNA-seq studies on the *Drosophila* eye represent a landmark in genomics. The application of single-cell RNA sequencing to the *Drosophila* eye not only showcases the versatility of this technology but also sets a new standard for cellular resolution in complex tissues. The meticulous identification of novel marker genes is particularly impressive, as it opens up new avenues for research and potential therapeutic targets. The broader implications of this research are particularly exciting. The ability to adapt these methodologies to other model organisms and tissues holds great promise for advancing our understanding of cellular diversity and gene regulation across various biological systems. The potential for future research building on these findings is immense and could lead to significant breakthroughs in developmental biology and beyond. Overall, single-cell RNA-seq research is a testament to the power of innovation and meticulous research in unraveling the complexities of life at the cellular level. It not only contributes valuable insights to the scientific community but also inspires further exploration and discovery in the field of genomics.

Compliance with ethical standards

Disclosure of conflict of interest

The authors declare no conflicting interests.

Authors' Contributions

The manuscript was prepared by B.P.G and K.K.B.R. and reviewed by K.K.B.R.

References

- [1] Wawersik S, Maas RL: Vertebrate eye development as modeled in *Drosophila*. *Human Molecular Genetics* 2000, 9(6):917-925.
- [2] Prüßing K, Voigt A, Schulz JB: *Drosophila melanogaster* as a model organism for Alzheimer's disease. *Molecular Neurodegeneration* 2013, 8(1):35.
- [3] Sang T-K, Jackson GR: *Drosophila* models of neurodegenerative disease. *NeuroRx* 2005, 2:438-446.
- [4] Jeibmann A, Paulus W: *Drosophila melanogaster* as a model organism of brain diseases. *International journal of molecular sciences* 2009, 10(2):407-440.
- [5] Yamaguchi M, Yoshida H: *Drosophila* as a Model Organism. In: *Drosophila Models for Human Diseases*. Edited by Yamaguchi M. Singapore: Springer Singapore; 2018: 1-10.
- [6] Tomlinson A, Ready DF: Neuronal differentiation in *Drosophila ommatidium*. *Dev Biol* 1987, 120(2):366-376.
- [7] Tang F, Barbacioru C, Wang Y, Nordman E, Lee C, Xu N, Wang X, Bodeau J, Tuch BB, Siddiqui A et al: mRNA-Seq whole-transcriptome analysis of a single cell. *Nature Methods* 2009, 6(5):377-382.
- [8] Kolodziejczyk AA, Kim JK, Svensson V, Marioni JC, Teichmann SA: The technology and biology of single-cell RNA sequencing. *Molecular cell* 2015, 58(4):610-620.
- [9] Ready DF, Hanson TE, Benzer S: Development of the *Drosophila* retina, a neurocrystalline lattice. *Dev Biol* 1976, 53(2):217-240.
- [10] Li H, Janssens J, De Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA, Brbić M, Spanier K et al: Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science* 2022, 375(6584):eabk2432.
- [11] Yeung K, Bollepogu Raja KK, Shim Y-K, Li Y, Chen R, Mardon G: Single cell RNA sequencing of the adult *Drosophila* eye reveals distinct clusters and novel marker genes for all major cell types. *Communications Biology* 2022, 5(1):1370.
- [12] Saliba A-E, Westermann AJ, Gorski SA, Vogel J: Single-cell RNA-seq: advances and future challenges. *Nucleic Acids Research* 2014, 42(14):8845-8860.
- [13] Dion WA, Shittu MO, Steenwinkel TE, Raja KKB, Kokate PP, Werner T: The modular expression patterns of three pigmentation genes prefigure unique abdominal morphologies seen among three *Drosophila* species. *Gene Expression Patterns* 2020, 38:119132.
- [14] Raja KK, Shittu MO, Nouhan PM, Steenwinkel TE, Bachman EA, Kokate PP, McQueeney A, Mundell EA, Armentrout AA, Nugent A: The regulation of a pigmentation gene in the formation of complex color patterns in *Drosophila* abdomens. *Plos one* 2022, 17(12):e0279061.
- [15] Lun AT, McCarthy DJ, Marioni JC: A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* 2016, 5:2122.
- [16] Bollepogu Raja KK, Yeung K, Shim Y-K, Li Y, Chen R, Mardon G: A single cell genomics atlas of the *Drosophila* larval eye reveals distinct photoreceptor developmental timelines. *Nature Communications* 2023, 14(1):7205.
- [17] Raja KKB, Yeung K, Li Y, Chen R, Mardon G: A single cell RNA sequence atlas of the early *Drosophila* larval eye. *BMC Genomics* 2024, 25(1):616.
- [18] Wagner A, Regev A, Yosef N: Revealing the vectors of cellular identity with single-cell genomics. *Nature Biotechnology* 2016, 34(11):1145-1160.
- [19] Bollepogu Raja KK, Yeung K, Shim Y-K, Mardon G: Integrative genomic analyses reveal putative cell type-specific targets of the *Drosophila* ets transcription factor Pointed. *BMC Genomics* 2024, 25(1):103.
- [20] Shittu M, Steenwinkel T, Dion W, Ostlund N, Raja K, Werner T: RNA in situ hybridization for detecting gene expression patterns in the abdomens and wings of *Drosophila* species. *Methods and protocols* 2021, 4(1):20.
- [21] Raja KK, Bachman EA, Fernholz CE, Trine DS, Hobmeier RE, Maki NJ, Massoglia TJ, Werner T: The Genetic Mechanisms Underlying the Concerted Expression of the yellow and tan Genes in Complex Patterns on the Abdomen and Wings of *Drosophila guttifera*. *Genes* 2023, 14(2):304.
- [22] Gudelly BP, Raja KKB: Protocol for dissection of *Drosophila* abdomens for fluorescent imaging. *Int J Sci Res Arch* 2024, 12(2):1211-1216.

- [23] Bollepogu Raja T: The role of toolkit genes in the evolution of complex abdominal color patterns in *Drosophila guttifera*. 2017.
- [24] Trapnell C: Defining cell types and states with single-cell genomics. *Genome Res* 2015, 25(10):1491-1498.
- [25] Finet C, Kassner VA, Carvalho AB, Chung H, Day JP, Day S, Delaney EK, De Ré FC, Dufour HD, Dupim E et al: DrosoPhyla: Resources for Drosophilid Phylogeny and Systematics. *Genome Biology and Evolution* 2021, 13(8).
- [26] Liu L, Liu C, Quintero A, Wu L, Yuan Y, Wang M, Cheng M, Leng L, Xu L, Dong G et al: Deconvolution of single-cell multi-omics layers reveals regulatory heterogeneity. *Nature Communications* 2019, 10(1):470.